



UNIVERSITI PUTRA MALAYSIA

**DIGESTIBILITY OF INGREDIENTS IN TWO PELLETED DIETS BY
Macrobrachium rosenbergii (de Man)**

KENNETH CHIN SUI SIAN

FPSS 1988 2

DIGESTIBILITY OF INGREDIENTS IN TWO PELLETTED
DIETS BY Macrobrachium rosenbergii (de Man)

KENNETH CHIN SUI SIAN

MASTER OF SCIENCE

(FISHERIES)

UNIVERSITI PERTANIAN MALAYSIA



DIGESTIBILITY OF INGREDIENTS IN TWO PELLETED
DIETS BY Macrobrachium rosenbergii (de Man)

by

Kenneth Chin Sui Sian

A thesis submitted in partial fulfilment of the degree
of Master of Science in the Faculty of Fisheries
and Marine Science,
Universiti Pertanian Malaysia


February 1988

It is hereby certified that we have read this thesis entitled 'Digestibility of Ingredients in Two Pelleted Diets by Macrobrachium rosenbergii (de Man)' by Kenneth Chin Sui Sian, and in our opinion it is satisfactory in terms of scope, quality, and presentation as partial fulfilment of the requirements for the degree of Master of Science



.....
SULAIMAN M. YASSIN, Ph.D.

Professor/Dean of Graduate Studies
Universiti Pertanian Malaysia
(Chairman Board of Examiners)



.....
CHO, C. YOUNG, Ph.D.

Department of Nutritional Science
University of Guelph
Guelph, Ontario, CANADA
(External Examiner)



.....
MOHD. SALLEH KAMARUDIN, M.S.

Department of Fisheries Biology and Aquaculture
Universiti Pertanian Malaysia
(Internal Examiner)



.....
LAW AH THEEM, Ph.D.

Associate Professor
Department of Fishing Technology and
Marine Science
Universiti Pertanian Malaysia
(Internal Examiner/Supervisor)

This thesis was submitted to the Senate of Universiti
Pertanian Malaysia and was accepted as partial fulfilment of the
requirements for the Degree of Master of Science.

Date: 21 JUL 1988

Sulaiman M. Yassin
.....
SULAIMAN M. YASSIN, Ph.D.
Professor and
Dean of Graduate Studies

DEDICATION

To my wife Theresa and our families through
whom God has blessed me.

ACKNOWLEDGEMENT

The author wishes to express his heartfelt gratitude to his supervisor Dr. Law Ah Theem and co-supervisors, Dr. Ang Kok Jee and En. Mohd. Salleh Kamarudin for their advice guidance and patience during the course of this project. Sincere appreciation is also extended to En. Zakaria Md. Sah, Mr. Ravinthar Veellu, Mr. Ng Chee Kiat, En. Ahmad Fauzi Mohd Wazir and Mr. Yap Chee for all their help.

This project was funded by EEC Project No. TSDA 287 (MAL) (H).



TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vii
LIST OF PLATES	ix
LIST OF APPENDICES	x
ABSTRACT	xii
CHAPTER 1 INTRODUCTION	1
Current Status of Prawn Nutrition Research in Malaysia	4
LITERATURE REVIEW	6
Techniques Used in Digestibility Studies in Fish Nutrition	6
Digestibility Studies in Prawns	11
a. Protein	11
b. Fat	14
c. Carbohydrate	16
d. Inorganic components	17
e. Energy	17
Nutritional Requirements of Prawns	18
a. Protein	18
b. Fat	20
c. Carbohydrate	20
d. Minerals	21
e. Vitamins	22
f. Energy	23
Feedstuffs for Aquaculture	23



	<u>Page</u>
a. Rice Bran	24
b. Wheat Pollard	24
c. Coconut Oilcake	25
d. Soyabean Meal	25
e. Palm Oil Kernel Cake	26
f. Fish Meal	27
g. Shrimp Meal	28
 CHAPTER 2 MATERIALS AND METHODS	 29
Feed Preparation	29
The Prawns, Culture Systems and Experimental Design	 31
Determination of Physico-chemical Properties of the Culture Water	 37
a. Dissolved Oxygen	37
b. Temperature	37
c. pH	37
d. Ammonium-nitrogen	37
e. Phenolphthalein Alkalinity	37
Proximate analysis	38
a. Crude Protein Determination	38
(Micro-kjeldahl method)	
b. Lipid Determination	41
c. Moisture Determination	42
d. Ash Determination	43
e. Chromic Oxide Determination	43

	<u>Page</u>
f. Energy Determination	44
g. Carbohydrate Determination	48
Equations for the Evaluation of Digestibility	48
Statistical Analysis	49
CHAPTER 3: RESULTS	50
Phyhsico-chemical parameters	50
Digestibility results	50
Apparent Digestibility of the Reference Diets	51
Diet A (30% protein)	51
a. Juveniles	51
b. Adults	62
Diets B (40% protein)	63
a. Juveniles	63
b. Adults	63
Apparent Digestibility of Diets A and B in Female Prawns with Three Different Stages of Gonad Maturation	64
CHAPTER 4: DISCUSSION	65
Apparent Digestibility of the Reference Diet	65
Digestibility in Relation to Gonad Maturation	76
Digestibility of the Ingredients	77
CHAPTER 5: CONCLUSIONS AND RECOMMENDATION	85
REFERENCES	88
APPENDICES	98

LIST OF TABLES

	<u>Page</u>
TABLE	
I Amounts of Feed Ingredients Used in Formulation of Diet A and B	30
II Reference and Test Diets used in this study	30
III Apparent Digestibility of Nutrients in the Ingredients of Diet A (30% protein) for Juvenile <u>M. rosenbergii</u>	52
IV Apparent Digestibility of Nutrients in the Ingredients of Diet B (40% protein) for Juvenile <u>M. rosenbergii</u>	53
V Apparent Digestibility of Nutrients in the Ingredients of Diet A (30% protein) for Adult <u>M. rosenbergii</u>	54
VI Apparent Digestibility of Nutrients in the Ingredients of Diet B (40% protein) for Adult <u>M. rosenbergii</u>	55
VII Apparent Digestion Coefficients of the Ingredients in Diet A in Juvenile <u>M.</u> <u>rosenbergii</u>	56
VIII Apparent Digestion Coefficients of the Ingredients in Diet A in Adult <u>M.</u> <u>rosenbergii</u>	57
IX Apparent Digestion Coefficients of the Ingredients in Diet B in Juvenile <u>M. rosenbergii</u>	58



	<u>Page</u>
TABLE	
X	Apparent Digestion Coefficients in the Ingredients in Diet B in Adult <u>M.</u> <u>rosenbergii</u> 59
XI	Apparent Digestibility of Nutrients in the Ingredients in Diet A (30% protein) in Female <u>M. rosenbergii</u> 60
XII	Apparent Digestibility of Nutrients in Diet B (40% protein) in Female <u>M.</u> <u>rosenbergii</u> 61
XIII	Summary of some Digestibility Deter- minations in Prawns and Fish 68



LIST OF PLATES

PLATE		<u>Page</u>
I	Culture Tanks for Juvenile Prawns	33
II	Cages Housing Juvenile Prawns	34
III	Culture Tanks for Adult Prawns	35
IV	Kjeldahl Digestion Flasks	39
V	Automatic Analyzer for Crude Protein Determination	40
VI	Double Beam Spectrophotometer	45
VII	Bomb Calorimeter	46



LIST OF APPENDICES

APPENDIX		<u>Page</u>
A	Comparison of Soxhlet and Microtechnique in Lipid Extraction	97
B	Standard for Chromic Oxide determination	98
C	Standard Curve of Absorbance against Concentration of Chromic Oxide (mg/100ml)	99
D	Proximate Chemical Analyses of Feed and Feces of the Reference Diet A and Test Diets in Juvenile <u>M. rosenbergii</u>	100
E	Proximate Chemical Analyses of Feed and Feces of the Reference Diet A and Test Diets in Adult <u>M. rosenbergii</u> ...	103
F	Proximate Chemical Analyses of Feed and Feces of the Reference Diet B and Test Diets in Juvenile <u>M. rosenbergii</u>	108
G	Proximate Chemical Analyses of Feed and Feces of the Reference Diet B and Test Diets in Adult <u>M. rosenbergii</u>	111
H	Proximate Chemical Analyses of Feed and Feces of the Reference Diet A and B in Female <u>M. rosenbergii</u> with Green, Yellow and Orange Gonads	114
I	Results of 1-way ANOVA test and Duncan's Multiple Range test for the Digestion Coefficients of the Ingredients of Diet A (30% protein) fed to Juvenile <u>M. rosenbergii</u>	115



APPENDIX

Page

J	Results of 1-way ANOVA Test and Duncan's Multiple Range Test for the Digestion Coefficients of the Ingredients of Diet B (40% protein) fed to Juvenile Prawns	116
K	Results of 1-way ANOVA Test and Duncan's Multiple Range Test for the Digestion Coefficients of the Ingredients of Diet A (30% protein) fed to Adult Prawns	117
L	Results of 1-way ANOVA Test and Duncan's Multiple Range Test for the Digestion Coefficients of the Ingredients of Diet B (40% protein) fed to Adult Prawns	118
M	Results of 1-way ANOVA Test of Ingredients of Diet A (30% protein) fed to Female Prawns	119
N	Results of 1-way ANOVA Test of Ingredients of Diet B (40% protein) fed to Female Prawns	120



ABSTRACT

An abstract of the thesis submitted to the Senate of Universiti Pertanian Malaysia as partial fulfilment of the requirements for the degree of Master of Science.

DIGESTIBILITY OF INGREDIENTS IN TWO PELLETTED DIETS BY Macrobrachium rosenbergii (de Man)

by

Kenneth Chin Sui Sian

1988

Chief Supervisor : Law Ah Theem, Ph.D.

Co-Supervisors : Ang Kok Jee, Ph. D.

Mohd Salleh Kamaruddin, M.S.

Faculty : Fisheries And Marine Science

The digestion coefficients of nutrients in two diets containing 30% and 40% protein were evaluated in the Malaysian giant freshwater prawn Macrobrachium rosenbergii. Chromic oxide was used as an internal marker for the evaluation of digestibility. Evaluations were carried out with juvenile and adult prawns including female prawns undergoing three different stages of gonad maturation.

The results indicate that the adult prawns were able to digest the nutrients of the ingredients better than the juveniles. In the two diets tested copra cake, wheat flour and soyabean meal were better digested than fish meal and shrimp meal for both juvenile and adult prawns. Copra cake was found to be a good source of lipids (71-100%), carbohydrate (85-96%) and dry matter (72-92%) as well as a good attractant. Carbohydrate, gross energy and dry matter were very well digested in wheat flour with values of 93-100%, 86-95% and 85-97% respectively. Soyabean meal gave good digestion coefficients for protein (84-99%), lipid (90-97%) and carbohydrate (80-100%). The results indicated that copra cake, wheat flour and soyabean meal are good sources of nutrients for M. rosenbergii. No differences in digestibility was observed for the female prawns with green, yellow and orange gonads. This showed that the stage of gonad maturation has no significant effect on digestibility.



ABSTRAK

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi sebahagian dari keperluan untuk Ijazah Master Sains.

KEBOLEHCERNAAN BAHAN-BAHAN DALAM DUA MAKANAN BERUNTIL OLEH Macrobrachium rosenbergii (de Man).

Oleh

Kenneth Chin Sui Sian

1988

Ketua Penyelia : Law Ah Theem, Ph.D.

Penyelia bersama : Ang Kok Jee, Ph. D.
Mohd Salleh Kamaruddin, M.Sc.

Fakulti : Perikanan Dan Sains Samudra

Kebolehcernaan nutrien dalam dua makanan yang mengandungi 30% dan 40% protein telah dikaji dalam udang Macrobrachium rosenbergii. Kromik oksida telah digunakan sebagai bahan penanda. Kajian ini telah menggunakan udang juvenil, dan udang dewasa serta udang betina yang berada dalam tiga tahap kematangan gonad yang berlainan.

Keputusan kajian ini menunjukkan bahawa udang dewasa lebih cekap menghadam nutrien bahan-bahan makanan daripada udang juvenil. Dari kedua-dua makanan beruntill yang dikaji didapati bahawa kelapa, gandum dan kacang soya lebih cekap dicerna dari serbuk udang atau serbuk ikan. Kelapa merupakan sumber yang baik untuk lemak (71-100%), karbohidrat (85-96%) dan bahan kering (72-92%). Ia juga merupakan bahan penarik yang baik untuk udang ini. Karbohidrat, tenaga kasar dan bahan kering dicerna dengan cekap dalam gandum dan nilai-nilainya adalah 93-100%, 86-95% dan 85-97%. Kacang soya telah memberi nilai-nilai yang tinggi bagi protein (84-99%), lemak (90-97%) dan karbohidrat (80-100%). Maklumat-maklumat ini menunjukkan bahawa kelapa, gandum dan kacang soya adalah sumber-sumber nutrien yang baik bagi M. rosenbergii. Bagi udang betina yang mempunyai gonad yang berwarna hijau, kuning dan oren tiada perbezaan dalam kebolehcernaan didapati. Ini bermakna tahap kematangan gonad tidak ada kesan terhadap kebolehcernaan.

CHAPTER 1

INTRODUCTION

The Malaysian giant freshwater prawn, Macrobrachium rosenbergii, is a high priced delicacy in Malaysia. It remains as an important candidate for freshwater aquaculture since there is a downtrend in its supply from capture fisheries due to decreasing wild stocks (Rabanal, 1982; Ong and Pang, 1982).

The culture of the giant freshwater prawn has been shown to be economically profitable in Taiwan (Liao and Chao, 1982), and in Malaysia (Tiensongrusme, 1983). In Malaysia prawn farming is fast expanding especially with regards to the marine prawn Penaeus monodon. However the mass production of the giant freshwater prawn has been hindered by some serious problems such as unreliable fry supply, poor management, and a lack of cost effective feeds. Although in Malaysia advances have been made in the mass production of M. rosenbergii postlarvae (Ang and Cheah, 1983), the development of a commercial feed for this species of prawn still lags behind.



In the natural environment, M. rosenbergii juveniles and adults have been reported to be omnivorous. Their diet consists mainly of various plants and animal material including detritus, grass roots, insect larvae, seeds, flesh and offal of fish (Ling and Merican, 1961; Ling, 1962; Fujimura, 1972). Prawns are deliberate feeders consuming large quantities of food and engage in frequent feeding activity due to its relatively short digestive tract. Food is located mainly by touch and smell with the use of the antennae and antennules. Prawns masticate their food ingesting small particles whole.

Traditional prawn culture in Malaysia has long relied on the natural productivity of the pond and to a lesser extent on supplemental feeds such as unprocessed agricultural wastes (copra cake, rice bran), grains, trash fish and commercial poultry pellets. However, the use of such materials have limitations especially due to handling and storage problems as well as the imbalance of nutrients. In any semi-intensive and intensive prawn culture system a nutritionally balanced diet has to be supplied in order for the prawn to grow at an optimal rate. It has been shown that substantial increase in the growth of juvenile M. rosenbergii can be obtained using formula feeds (Stahl, 1979, Fair and Fortner, 1981). Feed, however, accounts for a large portion of the total operational costs of a prawn farm. In South Carolina it was 42% of total costs (Roberts

and Bauer, 1978) and in Taiwan it was at least 30% of total costs (Liao and Chao, 1982). The costs of prawn feed varies considerably in South East Asia and in general it is expensive. In Malaysia, the price range is between 0.88 - 1.40 US\$/kg (Akiyama, 1987). The high cost of this feed is due to several reasons. The inadequate knowledge of prawn nutrition and high cost of feed ingredients are some of the reasons. There is also minimal or no use of least-cost programs in prawn feed formulations. This is due to the inability of the feed manufacturer to replace a feed ingredient with another to obtain the same feed quality. The ingredients used in the present feed are also very expensive because most of these are imported. There is a lack of information on the utilization of locally produced, non-conventional feed ingredients such as copra meal, rice bran and palm oil kernel. Since feed costs constitute a high proportion of the operating expenditure in prawn culture, feeds offer more opportunities for reducing production costs through their refinement. The development of an economically and nutritionally efficient prawn feed will not only cut costs but also improve production. Prawn feeds have been generally evaluated in terms of growth with less attention given to the digestibility of compounded diets or the individual feedstuffs by prawns. Digestibility data provide important nutritional information concerning feeds or feedstuffs that supplement growth measurements.

Current Status of Prawn Nutrition Research in Malaysia

The initial observations on the feeding habits of Macrobrachium rosenbergii was reported by Ling (1962). Poh (1985) simulated the amino acid profile of M. rosenbergii in test diets composed of fish meal, shrimp meal, soyabean meal, copra cake, and wheat flour. Linear programming was used to formulate diets with 25%, 30%, 40% and 50% protein levels. Growth trials experiments by Poh (1985) with postlarval and juvenile M. rosenbergii showed that the 40% diet gave the best growth whereas the 30% diet produced the best feed conversion ratio. From these results Poh recommended that the optimum protein level for young M. rosenbergii is 40%. Recent pond trials of M. rosenbergii under semi-intensive systems (Ang, 1987) at Universiti Pertanian Malaysia using Poh's formulation of 30% diet yielded an average of 873.2 kg/ha/158 days cycle and a FCR of 1.43 and 2.08.

Yong (1986) studied the effect of split rations on nursery production of M. rosenbergii. She reported that feeding 3 times a day resulted in a better survival than feeding once or twice a day.

Ang et. al. (1987) studied the effect of the binder, Basfin, on the growth of M. rosenbergii. They found that the optimum level of Basfin was 0.5%.

Cheah et al. (1987) studied the response of M. rosenbergii larvae development when fed a 40% protein diet (Poh, 1985), egg custard, cockles and M. lanchesterii flesh. They found that the larvae grew the best when fed with the 40% protein diet and Artemia.

Besides these studies, no systematic approach has been adopted to examine the prawns nutritional requirements with regard to locally available feedstuffs.

Although Poh's formulation produced the desired growth, the role of each component of the diet in terms of digestibility by the prawn was not understood. It has been shown that the absorption of nutrients by an organism will vary with the nutrient source. It is therefore necessary to obtain more information concerning the diet recommended by Poh particularly in terms of the digestion coefficients of the nutrients in each of the ingredients used. Hence this study was conducted to solve some of the problems.

The main objective of this study is to obtain the apparent digestibility and digestion coefficient values for dry matter and the nutrients in the 30% and 40% protein diets, and the ingredients used in the diets. It is hoped that the results of this study will enable the formulation of a feed which is cost-effective and which will promote better growth of the prawn.

LITERATURE REVIEW

Techniques Used In Digestibility Studies in Fish Nutrition

The development of an adequate feed in terms of nutritional content and digestibility is essential for the large scale production of M. rosenbergii. Not only must the diet contain ingredients of the proper nutritional quality and quantity, the nutrients of the ingredients must be easily absorbed. A feedstuff may appear from its chemical composition to be an excellent source of nutrients but will be of little actual value unless it can be digested and absorbed by the prawn. Lovell (1977) indicated that nutrient digestibility among feedstuffs varied considerably for different warmwater fishes. The same observation has been noted for prawns. For example, wheat starch was better digested than potato starch in the prawn, Palaemon serratus (Forster and Gabbott, 1971). Akiyama (1986) found that Penaeus vannamei digested soyabean proteins better than fish or shrimp proteins. In order to minimize food wastage, each dietary ingredient should be evaluated on the basis of digestibility. The information collected for various ingredients will provide a useful source of information concerning the types of ingredient suitable for use as prawn feeds. Knowledge of nutrient availability of the components of prawn feeds is desirable so that effective substitutions of one component for another may be achieved. Such

substitution can be important in the development of least-cost diets. However, the lack of knowledge of the nutrient availability of the different feedstuffs used in prawn feeds limits the attempts at ingredient substitution.

Digestibility is defined as the percentage of ingested nutrients which are not rejected as feces (Schneider and Flatt, 1975). The determination of digestibility involves measuring the amount of a specific nutrient or feedstuff ingested and subtracting that which is present in the feces following digestion. Endogenous material such as secretions from within the intestinal tract, sloughed epithelial cells, and other material of metabolic origin may also occur in the feces. Since most studies fail to consider such endogenous materials, apparent rather than true digestibility is reflected.

Digestibility can be measured using one of three different techniques. These are the total collection for gravimetric measurement, the use of indigestible tracers and the use of radioisotopic markers.

Gravimetric measurement of digestibility involves the measurement by weight of total food intake and total fecal production. A known amount of feed is given to the prawn and the egested material is collected and weighed. The difference between the amount consumed and that recovered is

the amount assimilated. The problem with this technique is the difficulty in measuring the total amount of ingestion and defecation accurately.

Various compounds have been utilized as indigestible tracers in the determination of digestibility. These include chromic oxide, long chain fatty acids, metallic iron powder, radioisotopic markers, non-hydrolyzable organic or inorganic constituents and total ash. The two markers most commonly used for aquatic animals are chromic oxide and total ash. By mixing an indigestible marker into the feed or by measuring an endogenous marker, digestibility can be calculated by comparing the ratio of the marker in the feed and the feces.

Conover (1966) introduced an ash-ratio technique for the determination of absorption efficiency in aquatic animals. This technique assumed that only the organic components of a feed is significantly absorbed during the digestion process. The ash is then considered as an indigestible marker in a diet and percent digestibility could be calculated by comparing the ash content of food and feces. Although this technique has been applied to a wide range of crustaceans including M. rosenbergii (Clifford and Brick, 1979), the assumption that the ash component of a ration is not absorbed may not hold true in all cases.

